

Mantodea (mantid), Diptera (fly) and *Suncus murinus* (Musk Shrew); also 13 Hemiptera (bugs) and five microchiropteran bats (Table 1).

DISCUSSION

Although biomass studies formed no part of this investigation, it must be mentioned that it is ecologically significant as it elucidates the total food mass consumed. For example, though the Scarabaeidae topped numerically with 454 individuals, they are significantly less in biomass than 78 mice. A casual calculation using the known weights of specimens (Kumar, 1985) shows the following: 454 Scarabaeidae at the estimated mean weight of 0.57g each weigh 258.78g, whereas 78 mice at the estimated mean weight of 13.30g each weigh 1037.40g.

It is interesting to compare the prey spectrum of *Athene brama* between the agricultural fields of Andhra Pradesh studied by Kumar (1985) and this biome. In Andhra Pradesh fields no precedent was recorded of the Spotted Owllet feeding on snails, dragonflies, mantids, termites, bugs, ant lion larvae, moths, flies and hymenopterans; all these were consumed here in addition to very significant quantities of termites (19.88%) and Hymenoptera (21.06%). Hence it could be concluded that the Spotted Owllet has opportunistic feeding habits - its prey varying with locality and availability of food. This trait was recognized by Kumar (1985) and could be the reason for it being the most abundant and widespread species of owl on the Indian sub continent.

REFERENCES

- Ali, S. and S.D. Ripley (1969). *Handbook of the Birds of India and Pakistan*. 3: 299-302. Oxford University Press.
- Agrawal, V.C. (2000). Taxonomic studies on the Indian Muridae and Hystricidae (Mammalia: Rodentia). *Records of the Zoological Survey of India*, Miscellaneous Publication. Occasional Paper No. 180: 1-177
- Bates, P.J.J. and D.L. Harrison (1997). *Bats of Indian Subcontinent*. Harrison Zoological Museum Publication, Kent, 258pp.
- Borror, D. (1992). *An Introduction to the Study of Insects*. Harcourt Brace College Publishers, USA, 876pp.
- Errington, P.L. (1930). The pellet analysis method of raptor food habits study. *Condor* 32: 292-296.
- Heinrich, E. (1994). *Biology and Management of Rice Insects*, 780pp.
- Kumar, T.S. (1985). The Life History of the Spotted Owllet (*Athene brama brama* Temminck) in Andhra Pradesh. Monograph of the Raptor Research Centre, Hyderabad, 241pp.
- Mani, M. (1990). *General Entomology*. Oxford & IBH Publishing, New Delhi, 912pp.
- Mason, C.W. and H.M. Lefroy (1912). The food of birds in India. *Memoirs of the Department of Agriculture of India, Pusa: Entomology Series*. Vol 3. 376pp.
- Verzhutskii, B. and M.E. Ramanujam (2002). On the Prey of the Collared Scops Owl *Otus bakkamoena* (Pennant) at Auroville, Pondicherry. *Zoos' Print Journal* 17(1): 939-940.

ACKNOWLEDGEMENT

We sincerely wish to thank the European Commission's Tropical Dry Evergreen Forest (ECTDEF) Project, Pitchandikulam Bio-Resource Centre and Shakti Herbarium for providing the means to carry out this and other studies on the fauna of the Kaliveli bioregion.



THE PREY OF THE GREATER FALSE VAMPIRE BAT *MEGADERMA LYRA* E. GEOFFROY AT KALIVELI, TAMIL NADU

Mario Eric Ramanujam¹ and Boris Verzhutskii²

¹Principal Investigator (Fauna), ECTDEF Project, Pitchandikulam Bio-Resource Centre and Research Associate, Gratitude Avian Rehabilitation, Auroville, Pondicherry 605101, India

²Entomologist, Palmyra Project, Auroville, Pondicherry 605111, India
Email: ¹ tdef@auroville.org.in; ² palmyra@auroville.org.in

The prey of *Megaderma lyra* has been well documented in general (Bates & Harrison, 1997; Hill & Smith, 1984; Prater, 1993), as well as in certain areas, e.g., in Rajasthan (Advani, 1981; Prakash, 1959), central and western India (Brosset, 1962), Bihar (Sinha, 1986) and Bangladesh (Khan, 2001). This report concerns the prey spectrum of this species in the Kaliveli area of Tamil Nadu in southern India.

Kaliveli (12°05'-12°15'N & 79°47'-79°59'E) is dominated by a 7.040ha lake, that is said to be one of the two most important wetlands along the Coromandel Coast of southern India (Perennou, 1987, 1989; Perennou & Santharam, 1990). Most of the surrounding land is taken up for farming, though a marginal fringe of the once widespread tropical dry evergreen forest survives in the form of sacred groves and fragmented forest blocks. Villages dot the entire region, and, in one such village - Nadukuppam (12°10'N & 79°53'E) a colony of over a hundred *Megaderma lyra* was discovered in a disused building. The entire floor was covered with faecal matter up to a couple of centimetres deep and dead vertebrate remains littered the entire place. On 3.xii.2003 a random sample of faeces was collected for analysis (represented in Table 1) and all vertebrate remains were collected for identification (Table 2).

The percentage of prey items consumed was assessed according to a pre-established systematic that is in place for calculating trophic connections of fauna in this region (Verzhutskii & Ramanujam, 2002). This methodology recognizes four classes: basic food (>20%), constant food (5-20%), supplementary food (1-5%) and chance food (< 1%).

Diagnosis from faeces: 3,936 prey items were identified in the random sample. Of these Blattaria accounted for a staggering 1,919 specimens (48.75%), and along with Coleoptera (22.4%) comprised the basic food. Among the Coleoptera, the families Carabidae, Elateridae, Chrysomelidae, Scarabaeidae, Hydrophilidae and Curculionidae were identified. 456 Isoptera (11.58%) comprised the constant food. Supplementary food consisted of Araneae (1.62%), Orthoptera (2.18%), Diptera (2.31%), Hymenoptera (3.96%) and Lepidoptera (4.39%). Chance food included Mollusca (0.27%), Neuroptera (0.33%), Mantodea (0.35%), Odonata (0.38%), Vertebrata (0.48%) and Hemiptera (0.94%). Specific identification of vertebrates was not possible from faecal matter.

Manuscript 1165; Received 13 March 2004; Finally accepted 10 August 2004;
© Zoo Outreach Organisation; www.zoosprint.org

Diagnosis from vertebrate carcass remains: 297 carcasses were collected. Geckos *Hemidactylus spp.* dominated with 45.79%. Frogs *Polypedates maculatus* and *Euphlyctis spp.* accounted for 25.58% and 9.42% respectively. Mice *Mus spp.* comprised 8.75%, followed by bats (7.07%). Among the 21 bat carcasses, 13 *Megaderma lyra* (4.37%) were identified, but it is not clear whether these were members of the colony that had died of natural causes or had been cannibalised. A snake was also found. In general, mammals accounted for 17.17%, reptiles 47.81% and amphibians 35%. Neither fish nor birds were in evidence.

Megaderma lyra has been recorded feeding on fish (Advani, 1981; Prater, 1993) and birds (Advani, 1981; Brosset, 1962; Khan, 2001; Prater, 1993; Sinha, 1986). Here, both fish and birds were conspicuous by their absence, though both are plentiful in the study area. Even more striking was the quantity of cockroaches predated on by *M. lyra* (48.75%) and have never been recorded to form an important prey base in other regions. This shows a marked prey preference that varies considerably from the diet of the species in other regions.

REFERENCES

Advani, R. (1981). Seasonal fluctuations in the feeding ecology of the Indian False Vampire *Megaderma lyra lyra* (Chiroptera: Megadermatidae) in Rajasthan. *Zeitschrift saugtierk* 46(2): 90-93
 Bates, P.J.J. and D.L. Harrison (1997). *Bats of the Indian Subcontinent*. Harrison Zoological Museum Publication, Kent, England, 258pp.
 Brosset, A. (1962). The Bats of central and western India. Part II. *Journal of the Bombay Natural History Society* 59(2): 583-624.
 Hill, J.E. and J.D. Smith (1984). *Bats: a Natural History*. British Museum, 243pp.
 Khan, M.A.R. (2001). Status and distribution of Bats in Bangladesh with notes on their ecology. *Zoos' Print Journal* 16(5): 479-483.
 Perennou, C. (1987). Two important wetlands near Pondicherry. *Blackbuck* 3: 1-9
 Perennou, C. (1989). Southern wintering range of some waterbirds. *Journal of the Bombay Natural History Society* 86(2): 247-248.
 Perennou, C. and V. Santharam (1990). An anthropological survey of some wetlands in south-east India. *Journal of the Bombay Natural History Society* 87(2): 354-363.

Table 2. Vertebrate Prey of *Megaderma lyra* as diagnosed from carcass remains

Prey	Scientific name	#	%
Musk shrew	<i>Suncus murinus</i>	2	0.67
Leaf-nosed bats	<i>Hipposideros spp.</i>	7	2.35
Pipistrelle	<i>Pipistrellus sp.</i>	1	0.33
Greater False Vampire	<i>Megaderma lyra</i>	13	4.37
Mice	<i>Mus spp.</i>	26	8.75
House Rat	<i>Rattus rattus</i>	1	0.33
Soft-furred Field Rat	<i>Millardia meltada</i>	1	0.33
Geckos	<i>Hemidactylus spp.</i>	136	45.79
Garden Lizard	<i>Calotes versicolor</i>	5	1.68
Buff-striped Keelback	<i>Amphiesma stolata</i> (Snake)	1	0.33
Indian Tree Frog	<i>Polypedates maculatus</i>	76	25.58
Pond frogs	<i>Euphlyctis spp.</i>	28	9.42
Total		297	

Prakash, I. (1959). Food of the Indian False Vampire. *Journal of Mammalogy* 40(4): 545-546.
 Prater, S.H. (1993). *The Book of Indian Animals*. 4th impression. Bombay Natural History Society & Oxford University Press, 324pp.
 Sinha, Y.P. (1986). *The Bats of Bihar: Taxonomy and Field Ecology*. Records of the Zoological Survey of India. Occasional Paper No. 77, 60pp.
 Verzhutskii, B. and M.E. Ramanujam (2002). On the prey of the Collared Scops Owl (*Otus bakkamoena* (Pennant)) at Auroville, Pondicherry. *Zoos' Print Journal* 17(11): 939-940.

ACKNOWLEDGEMENTS

We sincerely wish to thank the European Commission's Tropical Dry Evergreen Forest Project, Pitchandikulam Bioresource Centre and Shakti Herbarium for supplying the materials and means to carry out faunal assessments in the Kaliveli Watershed region.



Table 1. The prey of *Megaderma lyra* as diagnosed from faecal matter

Prey	#	%	Category
Mollusca (Molluscs)	11	0.27	OF
Araneae (Spiders)	64	1.62	SF
Odonata (Dragonflies & Damselflies) 1	15	0.38	OF
Orthoptera (Grasshoppers, Crickets, etc.)	86	2.18	SF
Mantodea (Mantids)	14	0.35	OF
Blattaria (Cockroaches)	1919	48.75	BF
Isoptera (Termites)	456	11.58	CF
Hemiptera (Bugs)	37	0.94	OF
Neuroptera (Lacewings & Ant - lions)	13	0.33	OF
Coleoptera (Beetles)	882	22.40	BF
Diptera (Flies)	91	2.31	SF
Lepidoptera (Butterflies & Moths) 2	173	4.39	SF
Hymenoptera (Ants, Bees & Wasps) 3	156	3.96	SF
Vertebrata (Vertebrates)	19	0.48	OF
	3936		

- Number consumed; 1 - All Anisoptera (dragonflies); 2-16 Larvae, rest Ditrysia (moths); 3-113 Formicidae (ants); BF - Basic food (>20%); CF - constant food (5-20%); SF - supplementary food (1-5%); OF - chance food (<1%)